

SUPPLEMENTARY TABLE. Frequency of symptoms lasting >4 weeks among adults aged ≥18 years tested for SARS-CoV-2 since January 2020, who initially reported these symptoms, by test status — Porter Novelli Internet survey, United States, April 2021

Symptom	Test-positive*	n/Total	Test-negative†	n/Total
	respondents weighted % (95% CI)		respondents weighted % (95% CI)	
Any symptom [§]	76.2 (72.3–80.0)	465/603	69.6 (67.1–72.1)	1,058/1,526
1 symptom only	31.5 (27.5–35.5)	206/603	30.3 (27.8–32.9)	479/1,526
2 symptoms	16.1 (12.9–19.4)	99/603	15.4 (13.4–17.4)	238/1,526
3 or more symptoms	28.5 (24.4–32.6)	160/603	23.9 (21.5–26.3)	341/1,526
Hair loss	58.3 (45.1–71.4)	39/66	51.2 (43.2–59.2)	98/188
Cognitive dysfunction [¶]	55.5 (45.6–65.3)	64/122	57.9 (51.8–64.1)	170/292
Shortness of breath/breathlessness	52.8 (45.1–60.4)	94/203	45.3 (38.7–51.9)	115/260
Post-exertional malaise**	49.6 (37.3–61.8)	45/84	53.2 (42.5–63.8)	52/103
Problems sleeping [§]	49.4 (41.0–57.7)	81/172	60.2 (56.1–64.2)	389/652
Fatigue/tired/weakness	48.4 (42.1–54.7)	141/301	49.7 (45.3–54.2)	283/570
Palpitations (heart racing or pounding)	47.9 (34.2–61.6)	29/65	35.9 (28.0–43.9)	61/165
Other symptom(s)	46.7 (20.9–72.5)	8/17	72.4 (54.7–90.0)	24/32
Change in smell or taste	46.4 (39.5–53.3)	120/247	44.0 (32.6–55.5)	41/93
Change in mood	40.1 (30.5–49.8)	48/123	48.2 (43.2–53.3)	235/464
Joint/muscle pain [§]	36.7 (29.1–44.3)	68/201	58.4 (53.7–63.2)	274/480
Cough	36.2 (29.9–42.5)	105/280	34.1 (28.8–39.5)	119/363
Chest pain/pressure	35.2 (26.6–43.9)	52/144	28.8 (21.5–36.2)	54/181
Stomach pain	34.6 (25.0–44.2)	40/114	38.3 (32.4–44.3)	121/321
Headache	31.1 (25.2–37.0)	93/298	35.2 (31.3–39.1)	247/698
Diarrhea	24.6 (16.5–32.8)	32/144	26.0 (20.7–31.3)	81/315
Nausea/vomiting	23.7 (15.0–32.4)	28/112	22.6 (16.2–29.0)	47/203
Sore throat	16.9 (10.9–22.8)	31/194	16.4 (11.6–21.3)	43/251
Fever or chills [§]	12.4 (7.9–17.0)	34/270	22.7 (16.7–28.6)	51/210

Abbreviations: n = unweighted number with positive or negative SARS-CoV-2 test result; CI = confidence interval.

*Persons with at least one positive SARS-CoV-2 test result.

† Persons with all negative SARS-CoV-2 test results.

[§] p-value for weighted Wald chi-square test <0.05, indicating significant differences between those testing positive and those testing negative for SARS-CoV-2

[¶] Difficulty thinking clearly, concentrating, forgetfulness, memory loss, or 'brain fog'

** Worsening of symptoms after even minor physical, mental, or emotional exertion